

Stress levels, eating habits, and body weight of adults in Higher Education Institutions

Gabriela Baptista¹  Joana Pereira de Carvalho-Ferreira¹  Daniela de Assumpção²  Marina Padovan¹ 
Alice Helena de Souza Paulino¹  Caroline Dário Capitani¹ 

¹Laboratório Multidisciplinar em Alimentos e Saúde, Faculdade de Ciências Aplicadas, Universidade Estadual de Campinas – LabMAS/UNICAMP. Campinas/SP, Brasil.

²Programa de Pós-Graduação em Gerontologia, Faculdade de Ciências Médicas, Universidade Estadual de Campinas – UNICAMP. Campinas/SP, Brasil.

E-mail: alicehsp@unicamp.br

Graphical Abstract

Highlights

- More than one-third of students and education professionals exhibited high stress levels;
- Higher stress levels were associated with increased consumption of foods with poor nutritional quality;
- Adults with high stress levels presented greater fluctuations in body weight.



Abstract

To analyze the prevalence of perceived stress during the period of social isolation imposed by the Covid-19 pandemic, according to sociodemographic characteristics, food consumption, and other health-related behaviors, among adult workers and students from Higher Education Institutions in the state of São Paulo, Brazil. This is a cross-sectional study conducted in 2020 with 524 individuals. Participants completed an online questionnaire on perceived stress (Perceived Stress Scale), in addition to reporting the weekly frequency of consumption of healthy and unhealthy foods. Adjusted prevalence ratios (PR) and corresponding 95% confidence intervals were estimated using Poisson regression. The highest level of perceived stress reached 31.1% of the sample and was even higher among individuals with lower household income (≤ 3 minimum wages, PR=2.14; 95% CI: 1.22–3.73), those who consumed fried foods (PR=1.80; 95% CI: 1.15–2.82), those who consumed sweets ≥ 2 days per week (PR=1.69; 95% CI: 1.11–2.56), those who reported increased alcohol consumption (PR=1.76; 95% CI: 1.15–2.68), and those who reported body weight reduction (PR=2.62; 95% CI: 1.25–5.51) during the Covid-19 pandemic. The findings of this study show that more than one-third of students and education professionals exhibited high levels of stress, along with greater consumption of foods with low nutritional quality, increased alcohol use, and weight loss during Covid-19-related isolation.

Keywords: Ultra-processed Foods. Social Isolation. Anxiety. Body Weight Changes.

Associate Editor: Edison Barbieri
Mundo Saúde. 2025;49:e17412025
O Mundo da Saúde, São Paulo, SP, Brasil.
<https://revistamundodasaude.emnuvens.com.br>

Received: 18 march 2025.
Accepted: 26 august 2025.
Published: 26 september 2025.

INTRODUCTION

In 2020, public health measures were implemented that affected the daily lives of populations worldwide^{1,2}. Social isolation negatively impacted daily activities, compromising social, economic, and psychological aspects, and leading to changes in diet and physical activity³, which contributed to weight gain in part of the population⁴.

A narrative review by Hossain and colleagues analyzed the epidemiology of mental health problems during 2020 and suggested that this period could result in depression, anxiety, stress, irrational anger, impulsivity, sleep disorders, and emotional disturbances⁵. Furthermore, a systematic review with meta-analysis including 14 studies indicated that approximately one in four adults exhibited significant stress due to changes resulting from imposed public health measures⁶.

Stress levels and emotional factors may be associated with the consumption of nutritionally unbalanced foods. In this context, the scenario of social isolation may be related to a greater search for foods considered as “comfort foods” (those with compositional characteristics capable of promoting emotional well-being, albeit momentarily), such as ultra-processed foods⁷. These are characterized by high content of fats, sodium, food additives, and su-

gars, and lower amounts of protein, dietary fiber, and micronutrients⁸. Changes in eating patterns, therefore, are due not only to limited access to foods during the isolation period but also to the psychological consequences of a confinement condition, such as stress, fear, and concerns⁹.

The immediate impact of confinement can be observed through variations in diet quality and population stress levels^{7,10}. Beyond stress levels, other factors directly related to emotional vulnerability may be affected when one or more individuals are deprived of social interaction¹⁰. Thus, it is important to evaluate which potential associations may be related to stress levels under such environmental conditions. Therefore, the hypothesis of this study is that individuals with high stress levels during the Covid-19 pandemic are more likely to adopt behavioral risk factors, such as unhealthy eating. In this regard, the aim of the present study was to analyze the prevalence of perceived stress during the first year of social isolation imposed by the Covid-19 pandemic, according to sociodemographic characteristics, food consumption, and other health-related behaviors in adult students and/or workers from Higher Education Institutions (HEIs) in the state of São Paulo, Brazil.

METHODS

Study Design and Population

This was a cross-sectional study conducted with students and/or workers from Higher Education Institutions (HEIs) in the state of São Paulo, between September and December 2020. The sample was selected by convenience, through recruitment in four HEIs, via invitations disseminated on social media (Facebook, WhatsApp, and Instagram) and by e-mail. Data were collected using a pre-coded questionnaire, developed on the Google Forms platform, based on questions retrieved or adapted from previously validated health surveys, such as ConVid¹¹, Pesquisa de Comportamentos, and the Health and Food Consumption Survey of Campinas, SP¹². The questionnaire was self-administered by the participants using a computer or mobile phone with internet access. Prior to its application, the instrument was pilot tested. The study was con-

ducted after approval by the Research Ethics Committee of the University of Campinas (UNICAMP) (#4.213.274). Participants answered the questionnaire after providing informed consent (Informed Consent Form – ICF).

Variables

Perceived stress was the dependent variable of the study and was assessed using the Perceived Stress Scale questionnaire, translated and validated for university students¹³. This instrument consists of 14 items addressing the perception of stressful experiences (reflecting negative feelings and the ability to cope with stress) during the past month. Responses followed a five-point Likert-type scale, with the categories: never, almost never, sometimes, fairly often, and very often¹³. The total score ranged from 0 to 40 points, with higher scores in-

dicating greater stress levels. The overall score was categorized into distribution tertiles: 0–14 (tertile 1), 15–21 (tertile 2), and 22–40 points (tertile 3). Individuals in the upper tertile (≥ 22 points) were classified as having a high level of stress. The upper tertile (≥ 22 points) was used to analyze possible associations of higher perceived stress levels during the pandemic with the following independent variables:

a) Sociodemographic: sex (male and female), age group (≤ 30 , 31–40, 41–50, 51–59, and ≥ 60 years), education (high school completed/undergraduate incomplete; and undergraduate completed), and household income (≤ 3 , > 3 to ≤ 5 , and > 5 minimum wages).

b) Eating habits: the question “Usually, during the pandemic, on how many days of the week do you usually eat these foods?” was used to assess frequency of consumption of healthy foods (rice, beans, eggs, beef, chicken and fish, vegetables, fruits, fruit juice, whole foods, and whole grains) and unhealthy foods (ready-made dishes, instant noodles, pizza and snacks, packaged savory snacks, sweets, candies, soft drinks, processed meats, and fried foods). Responses were grouped according to regular consumption of healthy foods (≥ 5 days/week) and unhealthy foods (≥ 2 days/week).

c) Health-related behaviors, body weight change, and eating practices: hours of sleep (< 8 and ≥ 8 hours/day), physical activity (does not practice, 1–2 times/week, and ≥ 3 times/week), alcohol consumption (no change, increased, decreased), smoking (yes and no), changes in body weight (no, yes, weight gain, weight loss), food preparation at home (yes and no), ordering food via delivery apps (yes and no), and number of daily meals (1–3 and ≥ 4).

Data Analysis on Stress

Descriptive analyses (absolute and relative frequencies) were used to characterize the study participants. Subsequently, the prevalences of high perceived stress were estimated, and associations with independent variables were verified using Pearson’s Chi-square test with a significance level of 5% ($p < 0.05$). Adjusted prevalence ratios (PR) by sex and age and their corresponding 95% confidence intervals (95% CI) were then estimated. A multiple Poisson regression model was developed to identify variables independently associated with high stress levels and selected variables. All variables with $p < 0.20$ in the bivariate analysis were included in the model, and only those with significance $< 5\%$ remained. Data analyses were performed using Stata software, version 15.1¹⁴.

RESULTS

A total of 524 adults from Higher Education Institutions in the state of São Paulo completed the questionnaire. The characteristics of the sample are presented in Table 1. The majority of participants were female (59.3%), aged 51–59 years (24.43%); some were currently enrolled in higher education (20.8%), while most had already completed their degree (79.2%).

For the entire study population, the prevalence of high perceived stress reached 31.3% (95% CI: 27.5–35.4) and was lower among individuals over 40 years of age compared to younger adults (up to 30 years). Regarding eating habits, higher prevalences of stress were observed among those who regularly consumed fried foods, sweets, and soft drinks (Table 1).

Table 1 - Prevalence and prevalence ratios of perceived stress according to sociodemographic and dietary intake variables among students and workers from Higher Education Institutions, São Paulo, Brazil, 2020 (n = 524).

Variables and categories	n (%)	Prevalence (%)	p-value ^a	Adjusted PR ^b by sex and age (95% CI ^c)
Sex				
			0.005	
Male	213 (40.6)	24.4		1
Female	311 (59.4)	36.0		1.34 (0.96–1.87)
Total	524	31.3		
Age group (years)				
			<0.001	
≤ 30	104 (19.8)	51.9		1
31–40	96 (18.3)	42.7		0.87 (0.57–1.30)
41–50	124 (23.7)	27.4		0.54 (0.35–0.84)
51–59	128 (24.4)	18.7		0.38 (0.23–0.61)
≥ 60	72 (13.7)	15.3		0.32 (0.16–0.61)
Education				
			<0.001	
Undergraduate completed	415 (79.2)	26.5		0.91 (0.45–1.82)
High school completed/undergraduate incomplete	109 (20.8)	49.5		1
Household income (minimum wages)				
			0.001	
≤ 3	31 (7.6)	54.8		1.58 (0.88–2.84)
> 3 to ≤ 5	99 (24.4)	30.3		1.11 (0.71–1.74)
> 5	276 (68.0)	22.8		1
Dietary intake				
Rice (≥ 5 days/week)	203 (38.7)	36.4	0.043	1.20 (0.88–1.64)
Beans (≥ 5 days/week)	116 (22.1)	40.5	0.015	1.22 (0.86–1.72)
Eggs (≥ 5 days/week)	58 (11.1)	44.8	0.018	1.40 (0.92–2.13)
Vegetables (≥ 5 days/week)	257 (49.0)	30.7	0.787	1.00 (0.73–1.37)
Fruits (≥ 5 days/week)	260 (49.6)	29.2	0.311	0.97 (0.71–1.33)
Fruit juice (≥ 5 days/week)	91 (17.4)	38.5	0.098	1.38 (0.94–2.00)
Whole foods (≥ 5 days/week)	106 (20.2)	33.0	0.669	1.05 (0.72–1.53)
Red meat (≥ 5 days/week)	37 (7.1)	48.6	0.018	1.42 (0.86–2.33)
Chicken (≥ 5 days/week)	24 (4.6)	62.5	0.001	1.64 (0.95–2.81)
Fish (≥ 5 days/week)	159 (30.3)	28.9	0.441	1.00 (0.71–1.41)
Processed meats (≥ 2 days/week)	247 (47.1)	35.6	0.044	1.21 (0.89–1.65)
Ready-to-eat meals (≥ 2 days/week)	94 (17.9)	44.7	0.002	1.41 (0.99–2.00)
Fried foods (≥ 2 days/week)	124 (23.7)	50.0	<0.001	1.71 (1.22–2.38)
Pizza (≥ 2 days/week)	210 (40.1)	40.0	<0.001	1.27 (0.92–1.75)
Instant noodles (≥ 2 days/week)	39 (7.4)	43.6	0.085	1.23 (0.74–2.03)
Desserts (≥ 2 days/week)	218 (41.6)	39.4	0.001	1.34 (0.98–1.83)
Sweets/candies (≥ 2 days/week)	267 (50.9)	40.4	<0.001	1.58 (1.14–2.20)
Soft drinks (≥ 2 days/week)	152 (29.0)	44.1	<0.001	1.54 (1.12–2.10)
Packaged savory snacks (≥ 2 days/week)	54 (10.3)	44.4	0.028	1.33 (0.86–2.05)

Notes: ^ap-value from Pearson's Chi-square test (**p < 0.05**). ^bPR: Prevalence Ratio. ^cCI: 95% Confidence Interval.

Data showed that the prevalence of stress was lower among individuals who practiced physical exercise three or more times per week (PR = 0.68; 95% CI: 0.47–0.97). On the other hand, the prevalence of stress was twice as high among those who

reported body weight changes during the period of social isolation: PR = 2.06 (95% CI: 1.16–3.63) for those who reported weight gain and PR = 2.20 (95% CI: 1.21–4.00) for those who reported weight loss.

Table 2 - Prevalence and prevalence ratios of perceived stress according to lifestyle variables, body weight change, and eating practices among students and workers from Higher Education Institutions, São Paulo, Brazil, 2020 (n = 524).

Variables and categories	n (%)	Prevalence (%)	P-value ^a	Adjusted PR ^b by sex and age (95% CI ^c)
Hours of sleep/day				
			0.009	
< 8	292 (55.9)	26.7		1
≥ 8	230 (44.1)	37.4		1.20 (0.87–1.65)
Physical activity practice				
			0.038	
None	173 (33.0)	37.0		1
1–2 times/week	144 (27.5)	33.3		0.92 (0.63–1.34)
≥ 3 times/week	207 (39.5)	25.1		0.68 (0.47–0.97)
No change				
			<0.001	
Increased	294 (56.1)	22.4		1
Decreased	163 (31.1)	42.3		1.38 (0.94–2.03)
Decreased	67 (12.8)	43.3		1.37 (0.83–2.26)
Smoking				
			0.831	
Non-smoker	487 (92.9)	31.4		1
Smoker	37 (7.1)	29.7		0.87 (0.47–1.61)
Body weight change				
			<0.001	
No change	97 (20.2)	14.4		1
Weight gain	246 (51.4)	33.3		2.05 (1.16–3.63)
Weight loss	136 (28.4)	37.5		2.20 (1.21–4.00)
Home food preparation				
			0.004	
Yes	210 (40.1)	38.6		1.04 (0.71–1.52)
No	313 (59.8)	26.5		1
Food delivery (≥ 1 day/week)				
			<0.001	
Yes	260 (52.2)	39.2		1.41 (0.98–2.03)
No	238 (47.8)	21.0		1
Number of meals/day				
			0.025	
1–3	464 (88.7)	29.7		1
≥ 4	59 (11.3)	44.1		1.37 (0.90–2.10)

Notes: ^ap-value from Pearson's Chi-square test (**p < 0.05**). ^bPR: Prevalence Ratio. ^cCI: 95% Confidence Interval.

The results of the multiple Poisson regression model are presented in Table 3. Higher prevalences of stress were observed among individuals with lower household income (≤ 3 vs. > 5 minimum wages), those who consumed fried foods and sweets/candies at least twice per

week, those who reported increased alcohol consumption, and those who reported body weight reduction. The variable sex was retained in the model for adjustment, considering the differences in stress prevalence between men and women in the bivariate analysis.

Table 3 - Multiple Poisson regression model and variables associated with perceived stress among students and workers from Higher Education Institutions, São Paulo, Brazil (n = 524).

Variables and categories	Adjusted PR ^a	P-value ^b	95% CI ^c
Sex			
Male	1		
Female	1.36	0.154	0.89–2.08
Household income (minimum wages)			
< 3	2.14	0.008	1.22–3.73
> 3 to ≤ 5	1.39	0.154	0.88–2.20
> 5	1		
Dietary intake			
Fried foods (≥ 2 days/week)	1.80	0.010	1.15–2.82
Sweets/candies (≥ 2 days/week)	1.69	0.014	1.11–2.56
Alcohol consumption			
No change	1		
Increased	1.76	0.009	1.15–2.68
Decreased	1.25	0.550	0.60–2.63
Body weight change			
No change	1		
Weight gain	1.75	0.133	0.84–3.61
Weight loss	2.62	0.011	1.25–5.51

Notes: ^aAdjusted PR: prevalence ratio adjusted for all variables in the table. ^bp-value from Wald test. ^cCI: 95% Confidence Interval.

DISCUSSION

The present study investigated the prevalence of perceived stress among students and workers from Higher Education Institutions in the state of São Paulo, according to sociodemographic characteristics, food consumption, and other health-related behaviors. A high prevalence of elevated stress levels was found among students and workers from these HEIs in 2020, especially among women. High stress levels were significantly greater among individuals with lower household income, those who regularly consumed fried foods and sweets, those who reported increased alcohol intake, and those who reported body weight loss during the period of social isolation.

The HEIs assessed in this study were state institutions that maintained remote activities on an emergency basis, without a planning or adaptation period. Therefore, it is worth noting that the women evaluated were required to continue their professional and academic activities remotely during social isolation and possibly accumulated household tasks, as previously described in the literature. Although no significant association was observed between stress and sleep duration in this study, the epidemiological scenario of the Covid-19 pandemic implied an overloaded routi-

ne and excessive responsibilities for women, which, in turn, may have led to changes in sleep patterns, contributing to higher stress levels¹⁵. This finding is consistent with a study that identified a positive association between high prevalence of poor sleep quality and stress¹⁶. Data from a web-based cross-sectional study with Brazilian adults and older adults showed that, during the pandemic, 49.8% of women reported the onset of sleep problems and 53.7% reported worsening of preexisting sleep problems, prevalences that were 35% and 41% higher, respectively, than those observed among men¹⁷.

Higher level of perceived stress among women—identified in the bivariate analysis of the present study—was also reported by other authors^{18,19}. These findings may be related to the possibility that women have a higher propensity to perceive stress, which may also be reflected in dietary intake³. During social isolation, perceived stress among mothers was related to loss of working hours, job loss, and decreased ability to pay bills and obtain childcare support³. It has also been shown that women were at greater risk of developing mental health problems during the pandemic, as most remained responsible for the majority of domestic

tasks, including caring for children and older adults²⁰. Indeed, the literature highlights sex differences, indicating that women appear to face a higher lifetime risk of developing affective disorders, with stress being an important environmental factor associated with this risk¹⁹. Explanatory hypotheses include a potentially greater reactivity of the hypothalamic-pituitary-adrenal axis, known as the stress axis, in women, as well as greater exposure to negative social support²¹.

Regarding the higher prevalence of perceived stress among participants with lower household income, it is important to note that economic hardship and pressures resulting from this condition may lead to greater psychological strain, increased risk of anxiety and depression, and overall impairment of socioemotional development²². During social isolation, individuals with lower family income were likely those at greater risk of deprivation of basic resources, leading to situations of insecurity, such as food insecurity, which is associated with mental health risks, including depression and high levels of stress²³.

The results of the present study indicated a higher prevalence of perceived stress among participants who regularly consumed fried foods and sweets (two or more times per week), as well as those who reported changes in body weight (weight loss). These findings are consistent with similar studies conducted in other countries^{3,24,25}, highlighting a possible association between poorer psychological well-being and the consumption of less nutritious foods, along with variations in body weight, among students and workers from HEIs.

The relationship between perceived stress and food consumption has gained attention, as stress has been identified as a relevant factor associated with worsening quality of life and health. Among university students and staff in Chile, an increase in the consumption of comfort foods was observed due to increased emotional distress during social isolation¹⁸. Similarly, in France, 50.3% of participants in one study reported a negative change in perceived diet quality, which they attributed mainly to dietary choices that favored comfort food consumption²⁵. An online cross-sectional study conducted with 197 mothers of children aged 5 to 11 years showed that most participants resorted to comfort foods as a strategy to cope with perceived stress related to the Covid-19 pandemic³.

A statistical association was also observed between stress and increased alcohol consumption, which was 76% higher among individuals with elevated stress levels. Increased alcohol consumption during social restrictions was reported by 17.6% of the Brazilian adult population (aged 18 years or older), according to data from the virtual health survey 'ConVid, Pesquisa de Comportamentos'²⁶. Data from two online sur-

veys conducted between March and April 2020 with adults living in the United States using social media, showed that the adjusted odds ratios for reporting increased alcohol intake were 1.41 among respondents with anxiety symptoms and 1.64 among those with depressive symptoms, compared to those without such symptoms²⁷. Alcohol is recognized as a central nervous system depressant, and its use is associated with more than 230 diseases and health conditions, including mental disorders, which were likely exacerbated by alcohol consumption during the period of social isolation, triggering or worsening depressive and anxiety episodes²⁸.

The results of this study showed that body weight loss was 2.6 times more frequent among individuals who reported high levels of stress. It is relevant to highlight that, in situations of weight loss, stress may be a determining factor for some people, leading to reduced nutrient and caloric intake or the skipping of meals, as possible adaptive responses to stress^{29,30}. The attempt to minimize emotional discomfort through eating during social isolation may result in maladaptive eating behaviors, leading to weight fluctuations^{31,32}. Thus, stress may be considered a relevant factor for the increased susceptibility to body weight variations among students and workers from HEIs in Brazil, particularly in the state of São Paulo.

Although no direct association was observed in the present study, it can be speculated that greater consumption of energy-dense foods at the expense of fresh foods may be one of the contributing factors to body weight change in individuals with high levels of perceived stress, especially if the individual does not engage in or has low frequency of physical activity³³. This hypothesis should be further explored in future studies. Physical activity is not only important as a preventive behavior against obesity³⁴, but it is also associated with better psychological outcomes and healthier eating behaviors³⁵.

Supporting the role of physical exercise as a protective factor for health³⁶, a lower prevalence of perceived stress was observed among participants who practiced physical activity three or more times per week. Although this result was not independently associated with stress in the multiple regression model, it can be mentioned that, in general, higher perceived stress may contribute to reduced physical activity or increased sedentary behavior³⁷. Therefore, as reported in different studies^{2,7,37,38}, it is important to understand the nature of the associations between perceived stress during the period of social isolation in the pandemic and health-related behaviors. This is due to the need to prevent the long-term impacts of the pandemic on individual health, as well as the impact of other environmental situations that are potential sources of stress.

It is important to emphasize that the data from the present study may serve as a starting point for reflection and for strengthening institutional policies aimed at caring for the academic community. Collectively, the findings help to understand the impact of stress on students and staff at HEIs in the state of São Paulo, fostering discussion on efficient psychological support systems and structured support mechanisms for students, faculty, and staff. This is particularly relevant given the profound transformation in work modalities after the coronavirus pandemic, with a significant and seemingly lasting increase in partially remote work formats. In this sense, the perception of support from supervisors and colleagues appears to be a fundamental factor not only for increased technical security but also for the overall well-being and satisfaction of workers³⁹. It is also worth noting that, in more extreme situations of social isolation, and considering the expansion of hybrid forms of remote work (combining remote and in-person work), encouraging physical activity among members of the academic community appears to be a fundamental protective measure for psychological well-being.

This study has limitations that should be considered. The cross-sectional design does not allow establishing causality in the associations observed, such as the relationship between stress and increased alcohol consumption. Therefore, it is difficult to make inferences about possible changes in perceived stress levels over

time. The decision to categorize perceived stress into distribution tertiles was adopted due to the absence of validated cut-off points for the population studied. However, this strategy has limitations, as cut-off points are derived from the sample itself, which compromises the generalizability of the results. In addition, the sample was recruited by convenience and concentrated in four HEIs in the state of São Paulo, which limits the external comparability of the findings. Nevertheless, data reported in the scientific literature indicate an increase in stress levels in the population, with a higher prevalence of psychological disorders resulting from social isolation⁴⁰. Thus, it can be suggested that there was an increase in perceived stress during this period in the analyzed population. Furthermore, all data were collected through a self-administered online questionnaire, which may entail certain limitations, such as lack of accuracy regarding body weight data. Pre-pandemic data were collected retrospectively, meaning that eating habits and physical activity levels referring to the period prior to social isolation may have been influenced by recall bias. Finally, it is not possible to rule out the occurrence of non-response bias, considering that individuals in more adverse situations may have been less likely to participate in the study. However, the absence of data on non-respondents, if present, does not allow evaluation of the direction and magnitude of this bias. Such an effect could have led to an underestimation of the effects found in this study.

CONCLUSION

The results of this study reveal that the prevalence of perceived stress among students and workers from Higher Education Institutions in the state of São Paulo was significantly high during the period of social isolation imposed by the Covid-19 pandemic, especially among individuals who reported lower household income, frequent consumption of fried foods and sweets during the week, increased alcohol intake, and body weight loss. Overall, the findings emphasize the importance of psychological support strategies and targeted

interventions, particularly for academic and professional populations that faced, and continue to face, drastic changes in their routines and work environments. It is suggested that future efforts to promote mental well-being should include measures to encourage regular physical activity, healthy eating and and provide adequate emotional support. Despite its limitations, the data reflect a trend consistent with the literature on the impact of stress during social isolation and indicate areas for future research and intervention.

CRedit author statement:

Conceptualization: Carvalho-Ferreira, J.P.; Capitani, C. D. Methodology: Baptista, G.; Carvalho-Ferreira, J.P.; Assumpção, D.; Capitani, C. D. Validation: Baptista, G.; Carvalho-Ferreira, J.P.; Capitani, C. D. Statistical analysis: Assumpção, D. Formal analysis: Assumpção, D. Investigation: Baptista, G. Resources: Capitani, C. D.; Assumpção, D. Writing-original draft preparation: Baptista, G. Writing-review and editing: Carvalho-Ferreira, J.P.; Assumpção, D.; Padovan, M.; Paulino, A. H. S.; Capitani, C. D. Visualization: Carvalho-Ferreira, J.P.; Assumpção, D.; Padovan, M.; Paulino, A. H. S.; Capitani, C. D. Supervision: Capitani, C. D. Project administration: Capitani, C. D.

All authors have read and agreed to the published version of the manuscript.

Acknowledgments

The authors would like to thank all voluntary participants in the study and CNPq for granting the undergraduate research scholarship (Call 2021/2022 – PIBIC/CNPq).

Ethical statement

This study was fully approved by the Ethics Committee of the University of Campinas (UNICAMP) (#4.213.274) on August 21, 2020.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

REFERENCES

1. Holmes EA, O'Connor RC, Perry VH, et al. Multidisciplinary research priorities for the COVID-19 pandemic: a call for action for mental health science. *Lancet Psychiatry*. 2020 Jun;7(6):547-560.
2. Malta DC, Szwarcwald CL, Andrade MC, et al. The COVID-19 pandemic and changes in lifestyle of Brazilian adults: a cross-sectional study, 2020. *Epidemiol Serv Saude* [Internet]. 2020 Oct-Dec. [accessed August 13, 2025];29(4). Available at: <https://www.scielo.br/j/rbepid/a/4xc9dNPPnXXNMtSYpJX3kqh/?lang=en&format=pdf>
3. Wang SD, Devjani S, Chillakanti M, Dunton GF, Mason TB. The COMET study: Examining the effects of COVID-19-related perceived stress on Los Angeles mothers' dysregulated eating behaviors, child feeding practices, and body mass index. *Appetite*. 2021 Aug 1;163:105209.
4. Heriseanu AI, Hay P, Corbit L, Touyz S. Grazing in adults with obesity and eating disorders: a systematic review of associated clinical features and meta-analysis of prevalence. *Clin Psychol Rev*. 2017 Dec;58:16-32.
5. Hossain MM, Tasnim S, Sultana A, et al. Epidemiology of mental health problems in COVID-19: a review. *F1000Res*. 2020 Jun 23;9:636.
6. Cooke JE, Eirich R, Racine N, Madigan S. Prevalence of posttraumatic and general psychological stress during COVID-19: a rapid review and meta-analysis. *Psychiatry Res*. 2020 Oct;292:113347.
7. Di Renzo L, Gualtieri P, Pivari F, et al. Eating habits and lifestyle changes during COVID-19 lockdown: an Italian survey. *J Transl Med*. 2020 Jun 8;18(1):229.
8. Touvier M, da Costa Louzada M L, Mozaffarian D, Baker P, Juul F, Srour B et al. Ultra-processed foods and cardiometabolic health: public health policies to reduce consumption cannot wait *BMJ* [Internet] 2023 Oct. [accessed August 13, 2025]; 383. Available at: <https://www.bmj.com/content/383/bmj-2023-075294>
9. Naja F, Hamadeh R. Nutrition amid the COVID-19 pandemic: a multi-level framework for action. *Eur J Clin Nutr*. 2020 Aug;74(8):1117-1121.
10. Carroll N, Sadowski A, Laila A, Hruska V, Nixon M, Ma DWL, et al. The impact of COVID-19 on health behavior, stress, financial and food security among middle to high income Canadian families with young children. *Nutrients*. 2020 Aug 7;12(8):2352.
11. Barros MBA, Lima MG, Malta DC, Szwarcwald CL, Azevedo RCS, Romero D, et al. ConVid – Pesquisa de Comportamentos pela Internet durante a pandemia de COVID-19 no Brasil: concepção e metodologia de aplicação. *Cad Saúde Pública* [Internet]. 2021 [acessado em 13 de Agosto, 2025];37(3). Disponível em: <https://www.scielo.org/article/csp/2021.v37n3/e00268320>
12. Carvalho SDL, Barros Filho AAC, Assumpção MBA, Berti de Azevedo D, Qualidade da dieta segundo a autoavaliação de adolescentes: resultados do ISACamp Nutri. *Ciênc. Saúde Coletiva*. 2020;25(11):4451-4461.
13. Dias ACR, Andrade AC, Oliveira MS. Validation of the perceived stress scale for Brazilian university students. *Psicologia: Teoria e Prática*. 2015;17(1):28-36.
14. StataCorp. Stata Statistical Software: Release 15. College Station, TX: StataCorp LLC; 2017.
15. Dos Santos Quaresma MV, Marques CG, Magalhães ACO, Dos Santos RVT. Emotional eating, binge eating, physical inactivity, and vespertine chronotype are negative predictors of dietary practices during COVID-19 social isolation: a cross-sectional study. *Nutrients*. 2021 Oct;13(10):3558.
16. Almojali AI, Almalki SA, Allothman AS, Masuadi EM, Alaqeel MK. The prevalence and association of stress with sleep quality among medical students. *J Epidemiol Glob Health*. 2017 Sep;7(3):169-174.
17. Barros MBA, Lima MG, Malta DC, Szwarcwald CL, Azevedo RCS, Romero D, et al. Relato de tristeza/depressão, nervosismo/ansiedade e problemas de sono na população adulta brasileira durante a pandemia de COVID 19. *Epidemiol Serv Saúde* [Internet]. 2020 [acessado em 13 de Agosto, 2025];29(4). Disponível em: http://scielo.iec.gov.br/scielo.php?script=sci_abstract&pid=S1679-49742020000400021&lng=pt&nrm=iso
18. Aslan I, Ochnik D, Çınar O. Exploring perceived stress among students in Turkey during the COVID-19 pandemic. *Int J Environ Res Public Health*. 2020 Dec 2;17(23):8961.
19. Ochnik D, Rogowska AM, Kurnierz C, et al. Mental health prevalence and predictors among university students in nine countries during the COVID-19 pandemic: a cross-national study. *Sci Rep*. 2021 Sep 20;11(1):18644.
20. Almeida M, Shrestha AD, Stojanac D, Miller LJ. The impact of the COVID-19 pandemic on women's mental health. *Arch Womens Ment Health*. 2020 Dec;23(6):741-748.
21. Christiansen DM, McCarthy MM, Seeman MV. Where sex meets gender: how sex and gender come together to cause sex differences in mental illness. *Front Psychiatry*. 2022 Jun 28;13:856436.
22. Matthews KA, Gallo LC. Psychological perspectives on pathways linking socioeconomic status and physical health. *Annu Rev Psychol*. 2011;62:501-30.
23. Martin MS, Maddocks E, Chen Y, Gilman SE, Colman I. Food insecurity and mental illness: disproportionate impacts in the context of perceived stress and social isolation. *Public Health*. 2016;132:86-91.
24. Salazar-Fernández C, Palet D, Haeger PA, Román Mella F. The perceived impact of COVID-19 on comfort food consumption over time: the mediational role of emotional distress. *Nutrients*. 2021 Jun 2;13(6):1910.
25. Sarda B, Delamaire C, Serry AJ, Ducrot P. Changes in home cooking and culinary practices among the French population during the COVID-19 lockdown. *Appetite*. 2022 Jan 1;168:105743.
26. Malta DC, Gomes CS, Souza PRB Jr, Szwarcwald CL, Barros MBA, Lima MG, et al. A pandemia da COVID-19 e as mudanças no estilo de vida dos brasileiros adultos: um estudo transversal, 2020. *Epidemiol Serv Saúde* [Internet]. 2020 [acessado em 13 de Agosto, 2025];29(4). Disponível em: http://scielo.iec.gov.br/scielo.php?script=sci_abstract&pid=S1679-49742020000400025&lng=pt&nrm=iso
27. Capasso A, Jones AM, Ali SH, Foreman J, Tozan Y, DiClemente RJ. Increased alcohol use during the COVID-19 pandemic: the effect of mental health and age in a cross-sectional sample of social media users in the U.S. *Prev Med*. 2021;145:106422.
28. Sanchez ZM, Sanches YP, Cogo-Moreira H, Valente JY, Martins CB, Andreoni S, et al. Consumo de álcool durante a pandemia da COVID-19: uma reflexão necessária para o enfrentamento da situação. *Cad Saúde Pública* [Internet]. 2020 [acessado em 13 de Agosto, 2025];36(10). Disponível em: <https://www.scielo.org/article/csp/2020.v36n10/e00124520/>
29. Schultchen D, Reichenberger J, Mittl T, et al. Bidirectional relationship of stress and affect with physical activity and healthy eating. *Br J Health Psychol*. 2019 May;24(2):315-333.
30. Ozamiz-Etxebarria N, Dosil-Santamaria M, Picaza-Gorrochategi M, Idoiaga-Mondragon N. Levels of stress, anxiety and depression in the first phase of the COVID-19 outbreak in a sample collected in northern Spain. *Cad Saude Publica* [Internet]. 2020 Apr 27. [accessed August 13,

2025];36(4). Available at: <https://pubmed.ncbi.nlm.nih.gov/32374806/>

31. Araiza AM, Lobel M. Stress and eating: definitions, findings, explanations, and implications. *Soc Personal Psychol Compass* [Internet]. 2018 Apr. [accessed August 13, 2025];12(4). Available at: <https://compass.onlinelibrary.wiley.com/doi/full/10.1111/spc3.12378>
32. Frayn M, Knäuper B. Emotional eating and weight in adults: a review. *Curr Psychol*. 2018;37(4):924-933.
33. Pellegrini M, Ponzio V, Rosato R, et al. Changes in weight and nutritional habits in adults with obesity during the “lockdown” period caused by the COVID-19 virus emergency. *Nutrients*. 2020 Jul 7;12(7):2016.
34. Heatherton TF. Neuroscience of self and self-regulation. *Annu Rev Psychol*. 2011;62:363-390.
35. Martinez-Avila WD, Sanchez-Delgado G, Acosta FM, et al. Eating behavior, physical activity and exercise training: a randomized controlled trial in young healthy adults. *Nutrients*. 2020 Nov 29;12(12):3685.
36. Peluso MA, Andrade LH. Physical activity and mental health: the association between exercise and mood. *Clinics (Sao Paulo)*. 2005;60(1):61-70.
37. Tomiyama AJ. Stress and obesity. *Annu Rev Psychol*. 2019 Jan 4;70:703-718.
38. Ng DM, Jeffery RW. Relationships between perceived stress and health behaviors in a sample of working adults. *Health Psychol*. 2003 Nov;22(6):638-642.
39. Capone V, Schettino G, Marino L, et al. The new normal of remote work: exploring individual and organizational factors affecting work-related outcomes and well-being in academia. *Front Psychol*. 2024 Feb 12;15:1340094.
40. Shuja KH, Aqeel M, Jaffar A, Ahmed A. COVID-19 pandemic and impending global mental health implications. *Psychiatr Danub*. 2020 Spring;32(1):32-35.

How to cite this article: Baptista, G., Carvalho-Ferreira, J.P., Assumpção, D., Padovan, M., Paulino, A.H.S., Capitani, C.D. (2025). Stress levels, eating habits, and body weight of adults in Higher Education Institutions. *O Mundo Da Saúde*, 49. <https://doi.org/10.15343/0104-7809.202549e17412025>. *Mundo Saúde*. 2025,49:e17412025.